arXiv:0903.3510v1 [math.DG] 20 Mar 2009

HYPERSURFACES M^n IN $\mathbb{S}^k \times \mathbb{H}^{n-k+1}$

DANIEL KOWALCZYK

Abstract. Let $\psi : M^n \to \mathbb{S}^k \times \mathbb{H}^{n-k+1}$ be an isometric immersion of codimension 1, then there exist symmetric $(1,1)$-tensors S and f, a tangent vector field U and a smooth function λ on M^n that satisfy the compatibility equations of $\mathbb{S}^k \times \mathbb{H}^{n-k+1}$. In this paper, we will deal with the converse problem: "Given a Riemannian manifold M^n with symmetric $(1,1)$-tensors S and f, tangent vector field U and smooth function λ satisfying the conditions mentioned above, can M^n then be isometrically immersed in $\mathbb{S}^k \times \mathbb{H}^{n-k+1}$ in such a way that (g, S, f, U, λ) is realized as the induced structure?".

1. Introduction

It is well known that the Gauss and Codazzi equations are necessary conditions for a Riemannian manifold M^n to be locally isometrically immersed as hypersurface into an arbitrary Riemannian manifold $\widetilde{M}^{n+1}$. In the case that $\widetilde{M}^{n+1}(c)$ is a space form, the Gauss and Codazzi equations can be written in terms of the metric of M^n and of the shape operator S, which are both intrinsically known(as soon S is known). Moreover in this case the Gauss and Codazzi equations are sufficient for an n-dimensional Riemannian manifold to be isometrically immersed into the space $\widetilde{M}^{n+1}(c)$ as hypersurface with given second fundamental form S. In [2], B. Daniel gave a necessary and sufficient condition for an n-dimensional Riemannian manifold to be isometrically immersed in the Riemannian product of a sphere and the real line or the Riemannian product of a hyperbolic space and the real line in terms of its first and second fundamental forms and of the projection of the vertical vector field ∂_t on its tangent space. He rewrote the Gauss and Codazzi equations for hypersurfaces immersed in $\mathbb{S}^n \times \mathbb{E}^1$ or $\mathbb{H}^n \times \mathbb{E}^1$ in terms of the metric of M^n, the shape operator S, the projection T of the vertical vector field ∂_t onto the tangent space of M^n and the normal component ν of ∂_t. The Gauss and Codazzi equations together with extra conditions on T and ν, which follow from the fact that ∂_t is parallel in $\mathbb{S}^n \times \mathbb{E}^1$ or $\mathbb{H}^n \times \mathbb{E}^1$, are necessary and sufficient conditions for a Riemannian manifold to be isometrically immersed into $\mathbb{S}^n \times \mathbb{E}^1$ or into $\mathbb{H}^n \times \mathbb{E}^1$. In this paper we will extend this result to the Riemannian product of the k-dimensional sphere and the $(n-k+1)$-dimensional hyperbolic space of opposite sectional curvature.

On an arbitrary Riemannian product $M_1 \times M_2$ of two Riemannian manifolds there exist a natural symmetric $(1,1)$-tensor F such that $F^2 = I$ $(F \neq \pm I)$ and $\widetilde{\nabla} F = 0$, where $\widetilde{\nabla}$ is the Levi-Civita connection of the Riemannian product. F is called the product structure of $M_1 \times M_2$. Let M^n be a hypersurface of the Riemannian product $M_1 \times M_2$ with product structure F. We can put

$$FX = fX + u(X)\xi,$$

$$F\xi = U + \lambda\xi,$$

where X is a tangent vector field on M^n, ξ a unit normal of M^n in $M_1 \times M_2$, f is a $(1,1)$-tensor on M, U is a tangent vector field on M^n, u is a 1-form on M^n and λ is a smooth function on M^n. Moreover there are some conditions on f, U, u and λ which follow from the fact that $F^2 = I$, F is symmetric and $\widetilde{\nabla} F = 0$. In the case of $\mathbb{S}^k \times \mathbb{H}^{n-k+1}$ one can rewrite the Gauss equation in terms of S and f and the Codazzi equation in terms of u, as follows:

$$R(X,Y)Z = (SX \wedge SY)Z + \frac{1}{2}\left(f(X \wedge Y)Z + (X \wedge Y)fZ\right),$$
$$(\nabla_X S)(Y) - (\nabla_Y S)(X) = \frac{1}{2}\left(u(X)Y - u(Y)X\right).$$

Our main aim is to proof the following:

Theorem. *Let (M^n, g) be a simply connected Riemannian manifold of dimension n. Let S and f be symmetric fields of operators on M^n, U a vector field on M^n and λ a smooth function on M^n such that $f^2X = X - g(U,X)U$, $fU = -\lambda U$ and $\|U\|^2 + \lambda^2 = 1$. Assume that the structure (g, S, f, U, λ) satisfies the Gauss and Codazzi equation for hypersurfaces in $\mathbb{S}^k \times \mathbb{H}^{n-k+1}$ and the following equations:*

$$(\nabla_X f)(Y) = g(U,Y)SX + g(SX,Y)U, \qquad \nabla_X U = \lambda SX - fSX, \qquad X[\lambda] = -2g(SX,U).$$

Then there exist an isometric immersion $\psi : M^n \to \mathbb{S}^k \times \mathbb{H}^{n-k+1}$ such that the shape operator with respect to the normal ξ associated to f is S and such that

$$F(\xi) = \psi_*(U) + \lambda\xi, \tag{1}$$
$$F(\psi_* X) = \psi_*(f(X)) + g(U,X)\xi \tag{2}$$

for every tangent vector field X on M^n and whereby F denotes the product structure of $\mathbb{S}^k \times \mathbb{H}^{n-k+1}$. Moreover the immersion is unique up to the global isometries of $\mathbb{S}^k \times \mathbb{H}^{n-k+1}$.

First we will recall some basic facts about the theory of submanifolds ([1]) and some facts about Riemannian products. Then we will apply these facts to the case of hypersurfaces in the Riemannian product of the k-dimensional sphere and the $(n-k+1)$-dimensional hyperbolic space. In the last section we will give the proof of the above stated theorem.

2. Preliminaries

2.1. **Submanifold Theory.** Let $\psi : (M^n, g) \to (\widetilde{M}^{n+m}, \widetilde{g})$ be an isometric immersions of codimension m. Denote by ∇ and $\widetilde{\nabla}$ the Levi-Civita connections of respectively (M^n, g) and $(\widetilde{M}^{n+m}, \widetilde{g})$. The tangent vector fields on M^n will be denoted by $X, Y, Z, \ldots$ and normal vector fields along M^n in $\widetilde{M}^{n+m}$ will be denoted by $\xi, \xi_1, \xi_2, \ldots$. The formulas of Gauss and Weingarten which concern the decomposition of $\widetilde{\nabla}_X Y$ and $\widetilde{\nabla}_X \xi$, respectively, into their tangent and normal components are given by

$$\widetilde{\nabla}_X Y = \nabla_X Y + h(X,Y),$$
$$\widetilde{\nabla}_X \xi = -S_\xi X + \nabla^\perp_X \xi,$$

whereby h denotes the second fundamental form, S_ξ denotes the shape operator of M^n in $\widetilde{M}^{n+m}$ associated to the normal ξ and $\nabla^\perp$ denotes the normal connection of M^n in $\widetilde{M}^{n+m}$. The second fundamental form h and the shape operator S_ξ are related by

$$\widetilde{g}(h(X,Y),\xi) = g(S_\xi X, Y).$$

The shape operator S_ξ at a point p of M^n is a symmetric linear endomorphism on the tangent space T_pM^n for every normal vector ξ, since the second fundamental form h is a normal-bundle-valued symmetric $(1,2)-$tensor field on M^n.

Let $R, \widetilde{R}$ and $R^\perp$ denote the Riemannian curvature tensors of $\nabla, \widetilde{\nabla}$ and $\nabla^\perp$, respectively, taken with sign convention $R(X,Y) = [\nabla_X, \nabla_Y] - \nabla_{[X,Y]}$. One can easily deduce, using the formulae of Gauss and Weingarten that

$$\widetilde{R}(X,Y)Z = R(X,Y)Z + A_{h(X,Z)}Y - A_{h(Y,Z)}X + (\bar{\nabla}_X h)(Y,Z) - (\bar{\nabla}_Y h)(X,Z),$$

where $\bar{\nabla}$ is the van der Waerden-Bortolotti connection of M^n in $\widetilde{M}^{n+m}$ defined by

$$(\bar{\nabla}_X h)(Y,Z) = \nabla^\perp_X h(Y,Z) - h(\nabla_X Y, Z) - h(Y, \nabla_X Z).$$

The decomposition of $\widetilde{R}(X,Y)Z$ into a tangential component and normal component yields the equations of Gauss and Codazzi:

$$\tan(\widetilde{R}(X,Y)Z) = R(X,Y)Z + A_{h(X,Z)}Y - A_{h(Y,Z)}X,$$
$$\operatorname{nor}(\widetilde{R}(X,Y)Z = (\bar{\nabla}_X h)(Y,Z) - (\bar{\nabla}_Y h)(X,Z).$$

It is also possible to deduce the following equation of Ricci, again using the formulae of Gauss and Weingarten:

$$\operatorname{nor}(\widetilde{R}(X,Y)\xi) = R^\perp(X,Y)\xi + h(X, A_\xi Y) - h(A_\xi X, Y).$$

In the following we will mostly consider isometric immersions $\psi : (M^n, g) \to (\widetilde{M}^{n+1}, \widetilde{g})$ of codimension 1, i.e. hypersurfaces. Let ξ be a local unit normal vector field of M^n in $\widetilde{M}^{n+1}$; the formulae of Gauss and Weingarten can be written as

$$\widetilde{\nabla}_X Y = \nabla_X Y + h(X,Y)\xi,$$
$$\widetilde{\nabla}_X \xi = -SX,$$

whereby h denotes the scalar valued second fundamental form of M^n in $\widetilde{M}^{n+1}$ and S denotes the shape operator of M^n in $\widetilde{M}^{n+1}$, such that $g(SX,Y) = h(X,Y)$. The equations of Gauss and Codazzi then reduce to

$$\tan(\widetilde{R}(X,Y)Z) = R(X,Y)Z - (SX \wedge SY)Z,$$
$$\tan(\widetilde{R}(X,Y)\xi) = (\nabla_X S)(Y) - (\nabla_Y S)(X),$$

whereby $\wedge$ associates to two tangent vectors $v, w \in T_pM^n$ the endomorphism $v \wedge w$ of T_pM^n defined by

$$(v \wedge w)z = g(w,z)v - g(v,z)w$$

and whereby ∇S denotes the covariant derivative of the shape operator, i.e. whereby

$$(\nabla_X S)(Y) = \nabla_X(SY) - S(\nabla_X Y).$$

2.2. **Riemannian products.** Let M_1 and M_2 be two differentiable manifolds of dimensions $p \geq 2$ and $q \geq 2$, respectively. Consider the product manifold $M_1 \times M_2$ with natural projections $\pi_1 : M_1 \times M_2 \to M_1$ and $\pi_2 : M_1 \times M_2 \to M_2$. The tangent space $T_{(p_1,p_2)}(M_1 \times M_2)$ of $M_1 \times M_2$ at (p_1,p_2) is isomorphic to the direct sum $T_{p_1}M_1 \bigoplus T_{p_2}M_2$ for every point (p_1,p_2) of $M_1 \times M_2$, i.e. we can identify the vector space $T_{(p_1,p_2)}(M_1 \times M_2)$ with the vector space $T_{p_1}M_1 \bigoplus T_{p_2}M_2$. At every point (p_1,p_2) one can define the linear endomorphism $F_{(p_1,p_2)} : T_{(p_1,p_2)}(M_1 \times M_2) \to T_{(p_1,p_2)}(M_1 \times M_2) : (v_1, v_2) \mapsto (v_1, -v_2)$. Since $F_{(p_1,p_2)}$ is thus defined at every point of (p_1,p_2), we can construct a field of endomorphisms of the tangent spaces to $M_1 \times M_2$ by

$$F(X_1, X_2) = (X_1, -X_2),$$

whereby $X_1 \in TM_1$ and $X_2 \in TM_2$. Notice that $F^2 = I$, whereby I denotes the identity transformation on $T(M_1 \times M_2)$. It is easy to see that a tangent vector v to $M_1 \times M_2$ at (p_1, p_2) lies in $T_{p_1}M_1$ if and only if $Fv = v$ and that a tangent vector w to $M_1 \times M_2$ at (p_1, p_2) lies in $T_{p_2}M_2$ if and only if $Fw = -w$.

If (M_1, g_1) and (M_2, g_2) are Riemannian manifolds, consider the product Riemannian metric $\widetilde{g}$ on $M_1 \times M_2$:

$$\widetilde{g}((X_1, X_2), (Y_1, Y_2)) = g_1(X_1, Y_1) + g_2(X_2, Y_2),$$

for all $X_1, Y_1 \in TM_1$ and for all $X_2, Y_2 \in TM_2$. A product manifold of Riemannian manifolds equipped with the product Riemannian metric is called a Riemannian product manifold. Notice that the subspaces $T_{p_1}M_1$ and $T_{p_2}M_2$ of $T_{(p_1,p_2)}(M_1 \times M_2)$ are orthogonal and that F is a symmetric $(1,1)$-tensor on $M_1 \times M_2$ with respect to $\widetilde{g}$. Denote by $\widetilde{\nabla}$ the Levi-Civita connection of $(M_1 \times M_2, \widetilde{g})$. It can be shown that

$$(\widetilde{\nabla}_X F)(Y) = 0, \tag{3}$$

i.e. that

$$\widetilde{\nabla}_X (FY) = F(\widetilde{\nabla}_X Y).$$

Consider an isometric immersion $\psi : (M^n, g) \to (M_1 \times M_2, \widetilde{g})$ of codimension 1 with unit normal ξ, i.e. a hypersurface M^n in $M_1 \times M_2$. Then, we can put

$$FX = fX + u(X)\xi, \tag{4}$$

$$F\xi = U + \lambda\xi, \tag{5}$$

whereby f is a $(1,1)$-tensor on M^n, u is a 1-form on M^n, U is a tangent vector field on M^n and λ is a smooth function on M^n.

Lemma 1. *Suppose that M^n is a hypersurface of a Riemannian product $M_1 \times M_2$ with unit normal ξ and for which f, u, U and λ are defined as above. Then*

1. *f is a symmetric $(1,1)$-tensor such that $f^2X = X - u(X)U$ and that $fU = -\lambda U$,*
2. *$u(X) = g(U, X)$, and*
3. *$g(U, U) + \lambda^2 = 1$.*

Proof. Using the fact that F is a symmetric $(1,1)$-tensor field on $M_1 \times M_2$ with respect to the product Riemannian metric $\widetilde{g}$, we obtain that f is a symmetric $(1,1)$-tensor field on M^n and that $u(X) = g(U, X)$ for every tangent vector field of M^n:

$$\begin{aligned} g(fX, Y) &= \widetilde{g}(FX, Y) = \widetilde{g}(X, FY) = g(X, fY), \\ u(X) &= \widetilde{g}(FX, \xi) = \widetilde{g}(X, F\xi) = g(U, X). \end{aligned}$$

Using the equation $F^2 = I$ and comparing the tangent and the normal components, we obtain the other relations:

$$\begin{aligned} X &= F^2X \\ &= F(fX + g(U, X)\xi) \\ &= f^2X + g(U, fX)\xi + g(U, X)(U + \lambda\xi) \end{aligned}$$

and

$$\begin{aligned} \xi &= F^2\xi \\ &= F(U + \lambda\xi) \\ &= fU + g(U, U)\xi + \lambda(U + \lambda\xi). \end{aligned}$$

□

Notice that the third equation can also be obtained from $fU = -\lambda U$, when $U \neq 0$ in every point p of M^n.

Lemma 2. *Let $\psi : M^n \to M_1 \times M_2$ be an isometric immersion of codimension 1 with local unit normal ξ in a Riemannian product and for which f, U and λ are defined as above. Then*

1. $(\nabla_X f)(Y) = u(Y)SX + g(SX,Y)U,$
2. $\nabla_X U = \lambda SX - fSX$
3. $X[\lambda] = -2u(SX).$

Proof. We will use equation (3) to determine the covariant derivatives of f, U and λ. We will therefore find a expression of $(\widetilde{\nabla}_X F)(Y)$ in terms of f, U and λ using the formulas of Gauss and Weingarten:

$$\begin{aligned}0 &= \widetilde{\nabla}_X FY - F(\widetilde{\nabla}_X Y)\\ &= \widetilde{\nabla}_X (fY + g(U,X)\xi) - F(\nabla_X Y + g(SX,Y)\xi), and\\ &= (\nabla_X f)Y - g(U,Y)SX - g(SX,Y)U + (g(\nabla_X U,Y) + g(fSX,Y) - \lambda g(SX,Y))\xi\end{aligned}$$

and

$$\begin{aligned}0 &= \widetilde{\nabla}_X F\xi - F(\widetilde{\nabla}_X \xi)\\ &= \widetilde{\nabla}_X (U + \lambda\xi) + F(SX)\\ &= \nabla_X U - \lambda SX + fSX + (X[\lambda] + 2g(U,SX))\xi.\end{aligned}$$

By comparing the tangent and normal components, we obtain the equations of Lemma 2. We notice that the third equation can also be obtained form the third equation of Lemma 1 and the second equation of Lemma 2 when $\|U\| \neq 1$ for every point p of M^n □

3. Isometric immersions into $\mathbb{S}^k \times \mathbb{H}^{n-k+1}$

Denote by $\mathbb{L}^{n+3} = (\mathbb{R}^{n+3}, \langle .,. \rangle = dx_1^2 + \cdots + dx_{n+2}^2 - dx_{n+3}^2)$ the Minkowski space of dimension $n+3$ and define $\mathbb{S}^k \times \mathbb{H}^{n-k+1}$ as the following submanifold of $\mathbb{L}^{n+3}$, equipped with the induced metric:

$$\mathbb{S}^k \times \mathbb{H}^{n-k+1} = \{(x_1, \ldots, x_{n+2}, x_{n+3}) | x_1^2 + \cdots + x_{k+1}^2 = 1,\ x_{k+2}^2 + \cdots + x_{n+2} - x_{n+3}^2 = -1,\ x_{n+3} > 0\}.$$

Then $\mathbb{S}^k \times \mathbb{H}^{n-k+1}$ is the Riemannian product of the k-dimensional sphere and the $(n-k+1)$-dimensional hyperbolic space. The Riemannian product $\mathbb{S}^k \times \mathbb{H}^{n-k+1}$ is a submanifold of $\mathbb{L}^{n+3}$ with orthonormal normal vectors $\xi_1 = (x_1, \ldots, x_{k+1}, 0, \ldots, 0)$ and $\xi_2 = (0, \ldots, 0, x_{k+2}, \ldots, x_{n+3})$. Using the formula of Gauss for isometric immersions into semi-Riemannian manifolds, we find that the Levi-Civita connection $\widetilde{\nabla}$ of $\mathbb{S}^k \times \mathbb{H}^{n-k+1}$ is given by the following expression in terms of the Levi-Civita connection D of $\mathbb{L}^{n+3}$ and the product structure F of $\mathbb{S}^k \times \mathbb{H}^{n-k+1}$:

$$\widetilde{\nabla}_X Y = D_X Y + \frac{1}{2}\langle X + FX, Y\rangle \xi_1 - \frac{1}{2}\langle X - FX, Y\rangle \xi_2. \tag{6}$$

Consequently, we obtain that the Riemann-Christoffel curvature tensor $\widetilde{R}$ is given by

$$\widetilde{R}(X,Y)Z = \frac{1}{2}\left(F(X \wedge Y)Z + (X \wedge Y)FZ\right). \tag{7}$$

Consider a hypersurface M^n of $\mathbb{S}^k \times \mathbb{H}^{n-k+1}$ with unit normal ξ and shape operator S. Let ∇ and R be the Levi-Civita connection and the Riemann-Christoffel curvature of M^n. Let f be a endomorphism on TM^n, u a 1-form, U a vector field tangent to M^n and λ a smooth function on

M^n defined as in section 2.2. Since the product structure F of $\mathbb{S}^k \times \mathbb{H}^{n-k+1}$ is symmetric and satisfies $F^2 = I$, we obtain the equations of Lemma 1 for f, u, U and λ. Using these notations and equation (7), we obtain that the equations of Gauss and Codazzi reduce to

$$R(X,Y)Z = (SX \wedge SY)Z + \frac{1}{2}\left(f(X \wedge Y)Z + (X \wedge Y)fZ\right), \tag{8}$$

$$(\nabla_X S)(Y) - (\nabla_Y S)(X) = \frac{1}{2}\left(u(X)Y - u(Y)X\right). \tag{9}$$

We also have that the product structure F of $\mathbb{S}^k \times \mathbb{H}^{n-k+1}$ is parallel and hence we also obtain the equations of Lemma 2 for f, U and λ. The equations of Lemmas 1 and 2 together with the equations of Gauss and Codazzi for hypersurfaces of $\mathbb{S}^k \times \mathbb{H}^{n-k+1}$ are called the compatibility equations of $\mathbb{S}^k \times \mathbb{H}^{n-k+1}$. In the following we will suppose that $f \neq \pm Id$, because otherwise we would have that $U = 0$, $\lambda = \pm 1$ and

$$R(X,Y)Z = (SX \wedge SY)Z \pm (X \wedge Y)Z,$$
$$(\nabla_X S)(Y) - (\nabla_Y S)(X) = 0,$$

and hence M^n would be an isometrically immersed into $\mathbb{S}^{n+1}$ or into $\mathbb{H}^{n+1}$.

Definition 1. Let S and f be symmetric fields of operators on M^n, U a vector field on M^n and λ a smooth function on M^n. We say that the structure (g, S, f, U, λ) of a Riemannian manifold (M^n, g) satisfies the compatibility equations of $\mathbb{S}^k \times \mathbb{H}^{n-k+1}$ if the following equations are satisfied:

$$g(fX,Y) = g(X,fY) \qquad f^2X = X - g(U,X)U \qquad fU = -\lambda U \qquad g(U,U) + \lambda^2 = 1, \tag{10}$$

$$R(X,Y)Z = (SX \wedge SY)Z + \frac{1}{2}\left((fX \wedge fY)Z + (X \wedge Y)fZ\right), \tag{11}$$

$$(\nabla_X S)(Y) - (\nabla_Y S)(X) = \frac{1}{2}\left(u(X)Y - u(Y)X\right), \tag{12}$$

$$(\nabla_X f)(Y) = u(Y)SX + g(SX,Y)U, \tag{13}$$

$$\nabla_X U = \lambda SX - fSX, \tag{14}$$

$$X[\lambda] = -2u(SX). \tag{15}$$

We will show now that an arbitrary Riemannian manifold (M^n, g) with structure (g, S, f, U, λ) that satisfies the compatibility equations of $\mathbb{S}^k \times \mathbb{H}^{n-k+1}$ can be isometrically immersed in $\mathbb{S}^k \times \mathbb{H}^{n-k+1}$ and moreover that the immersion is unique up to isometries of $\mathbb{S}^k \times \mathbb{H}^{n-k+1}$.

Theorem 1. *Let (M^n, g) be a simply connected Riemannian manifold of dimension n. Let S and f be symmetric fields of operators on M^n, U a vector field on M^n and λ a smooth function on M^n. Assume that (g, S, f, U, λ) satisfies the compatibility equations for $\mathbb{S}^k \times \mathbb{H}^{n-k+1}$. Then there exists an isometric immersion $\psi : M^n \to \mathbb{S}^k \times \mathbb{H}^{n-k+1}$ such that the shape operator with respect to the normal ξ associated to ψ is S and such that*

$$F(\xi) = \psi_*(U) + \lambda\xi, \tag{16}$$

$$F(\psi_* X) = \psi_*(f(X)) + g(U,X)\xi \tag{17}$$

for every tangent vector field X on M^n and whereby F is the product structure of $\mathbb{S}^k \times \mathbb{H}^{n-k+1}$. Moreover the immersion is unique up to a global isometries of $\mathbb{S}^k \times \mathbb{H}^{n-k+1}$.

4. Proof of the theorem

In the following we will give the proof of the theorem. We will use the techniques of [3] and [4] to proof the theorem. Let TM^n be the tangent bundle of M^n. Suppose $N = M^n \times \mathbb{R}^3$ is the trivial bundle equipped with the Minkowski metric g_{-1}. Denote by $B = TM^n \oplus_W N$ the orthogonal Whitney sum of TM^n and the trivial bundle N. The metric on B will be denoted by $\widetilde{g}$. Let ξ, ξ_1 and ξ_2 be an orthonormal frame in N such that $g_{-1}(\xi,\xi) = g_{-1}(\xi_1,\xi_1) = -g_{-1}(\xi_2,\xi_2) = 1$. We define a connection D in B by

$$D_X Y = \nabla_X Y + g(SX,Y)\xi - \frac{1}{2}g(X+fX,Y)\xi_1 + \frac{1}{2}g(X-fX,Y)\xi_2, \tag{18}$$

$$D_X \xi = -SX - \frac{1}{2}g(U,X)(\xi_1+\xi_2), \tag{19}$$

$$D_X \xi_1 = \frac{1}{2}(X + f(X) + g(U,X)\xi), \tag{20}$$

$$D_X \xi_2 = \frac{1}{2}(X - f(X) - g(U,X)\xi). \tag{21}$$

It is easy verified that the connection D on B is compatible with the metric $\widetilde{g}$ on B, i.e.

$$X\widetilde{g}(\eta_1,\eta_2) = \widetilde{g}(D_X\eta_1,\eta_2) + \widetilde{g}(\eta_1, D_X\eta_2)$$

for every tangent vector field X on M^n and $\eta_1, \eta_2 \in B$. The curvature tensor of B with connection D will be denoted by R^D. The curvature tensor $R^D : TM^n \times TM^n \times B \to B$ is a trilinear map over the module $C^\infty(M^n)$ of smooth functions on M^n defined by

$$R^D(X,Y)\eta = D_X D_Y \eta - D_Y D_X \eta - D_{[X,Y]}\eta.$$

We will show that the connection D on B is flat, i.e. $R^D = 0$.

Lemma 3. $R^D = 0$.

Proof. We will only calculate $R^D(X,Y)Z$ for arbitrary vector fields X, Y, Z on M^n. We will obtain that $R^D(X,Y)Z = 0$, because of the equations (11), (12) and (13). Using the definition of the connection D, we have

$$\begin{aligned}
D_X D_Y Z &= D_X(\nabla_Y Z + g(SY,Z)\xi - \frac{1}{2}(g(Y+fY,Z))\xi_1 + \frac{1}{2}(g(Y-fY,Z))\xi_2) \\
&= \nabla_X\nabla_Y Z + g(SX,\nabla_Y Z)\xi \\
&- \frac{1}{2}(g(X+fX,\nabla_Y Z))\xi_1 + \frac{1}{2}(g(X-fX,\nabla_Y Z))\xi_2 \\
&+ (g(\nabla_X SY,Z) + g(SY,\nabla_X Z))\xi + g(SY,Z)(-SX - \frac{1}{2}g(U,X)(\xi_1+\xi_2)) \\
&- \frac{1}{2}(g(\nabla_X Y,Z) + g(Y,\nabla_X Z) + g(\nabla_X fY,Z) + g(fY,\nabla_X Z))\xi_1 \\
&- \frac{1}{2}(g(Y+fY,Z))(\frac{1}{2}(X+fX+g(U,X)\xi)) \\
&+ \frac{1}{2}(g(\nabla_X Y,Z) + g(Y,\nabla_X Z) - g(\nabla_X fY,Z) - g(fY,\nabla_X Z))\xi_2 \\
&+ \frac{1}{2}(g(Y-fY,Z))(\frac{1}{2}(X-fX-g(U,X)\xi))
\end{aligned}$$

and a similar equation when X and Y are interchanged. We also have

$$\begin{aligned}D_{[X,Y]}Z &= \nabla_{[X,Y]}Z + g(S[X,Y],Z)\xi \\ &\qquad - \frac{1}{2}(g([X,Y]+f[X,Y],Z))\xi_1 + \frac{1}{2}(g([X,Y]-f[X,Y],Z))\xi_2.\end{aligned}$$

Hence we obtain that $R^D(X,Y)Z$ is given by

$$\begin{aligned}&R(X,Y)Z - (SX\wedge SY)Z - \frac{1}{2}(f(X\wedge Y)Z + (X\wedge Y)fZ) \\ &+ (g(\nabla_X SY,Z) - g(\nabla_Y SX,Z) - g(S[X,Y],Z) - \frac{1}{2}g(U,X)g(Y,Z) + \frac{1}{2}g(U,Y)g(X,Z))\xi \\ &- \frac{1}{2}(g(U,X)g(SY,Z) + g(\nabla_X fY,Z) - g(U,Y)g(SX,Z) - g(\nabla_Y fX,Z) - g(f[X,Y],Z))\xi_1 \\ &+ \frac{1}{2}(g(SX,Z)g(U,Y) - g(SY,Z)g(U,X) - g(\nabla_X fY,Z) + g(\nabla_Y fX,Z) + g(f[X,Y],Z))\xi_2.\end{aligned}$$

From equation (11), (12) and (13) we can conclude that $R^D(X,Y)Z$ vanishes for all tangent vector fields X, Y and Z. The cases $R^D(X,Y)\xi, R^D(X,Y)\xi_1$ and $R^D(X,Y)\xi_2$ can be treated analogously using equations (12), (13), (14) and (15). Since R^D is a trilinear map, we obtain that $R^D(X,Y)\eta = 0$ for every $\eta \in B$. ☐

We define now a bundle map F on B by

$$FX = fX + g(U,X)\xi, \tag{22}$$

$$F\xi = U + \lambda\xi, \tag{23}$$

$$F\xi_1 = \xi_1, \tag{24}$$

$$F\xi_2 = -\xi_2. \tag{25}$$

In the following two lemmas we will show that $F^2 = I$, F is symmetric with respect to $\widetilde{g}$ and the covariant derivative of the bundle map F on B is 0.

Lemma 4. *$F^2\eta = \eta$ and $\widetilde{g}(F\eta_1,\eta_2) = \widetilde{g}(\eta_1,F\eta_2)$.*

Proof. This follows immediately form the definition of the bundle map F and (10). ☐

Lemma 5. *$(D_X F)(\eta) = 0$ for every $X \in TM^n$ and every $\eta \in B$.*

Proof. This follows immediately form the definition of the bundle map F and equations (13), (14) , (15). ☐

Let B_1 and B_{-1} be subsets of B defined respectively by

$$\{\eta \in B \,|\, F\eta = \eta\}$$

and

$$\{\eta \in B \,|\, F\eta = -\eta\}.$$

In the next proposition we will show that there exist orthonormal parallel sections $\widetilde{\eta}_1,\dots,\widetilde{\eta}_{k+1}$, $\bar{\eta}_1,\dots,\bar{\eta}_{n-k+2}$ with $k \in \{1,\dots,n\}$ such that $F\widetilde{\eta}_i = \widetilde{\eta}_i$ and $F\bar{\eta}_\alpha = -\bar{\eta}_\alpha$ for $i \in \{1,\dots,k+1\}$ and $\alpha \in \{1,\dots,n-k+2\}$.

Proposition 1. *Let (M^n,g) be a Riemannian manifold with structure (g,S,f,U,λ) that satisfies the compatibility equations of $\mathbb{S}^k \times \mathbb{H}^{n-k+1}$. Let B be the vector bundle over M^n as defined above and F the bundle map of B as defined above, then there exist orthonormal parallel* *sections* $\widetilde{\eta}_1, \dots, \widetilde{\eta}_{k+1}, \bar{\eta}_1, \dots, \bar{\eta}_{n-k+2}$ *with* $k \in \{1, \dots, n\}$ *such that* $\widetilde{\eta}_1, \dots, \widetilde{\eta}_{k+1} \in B_1$ *and that* $\bar{\eta}_1, \dots, \bar{\eta}_{n-k+2} \in B_{-1}$.

Proof. Let p be a point of M^n such that $f_p \neq \pm Id$. By definition of the bundle map F, we obtain that Fv lies in the subspace V of B_p spanned by T_pM^n and ξ_p for every $v \in T_pM^n$. Analogously we have that $F\xi_p$ lies in V. Since F is symmetric with respect to $\widetilde{g}$, $\widetilde{g}$ is positive definite when restricted to V, $F^2 = Id$, $F(V) \subset V$ and $f_p \neq Id$, there exist an orthonormal basis $\{\widetilde{v}_1, \dots, \widetilde{v}_k, \bar{v}_1, \dots, \bar{v}_{n-k+1}\}$ with $k \in \{1, \dots, n\}$ such that $F\widetilde{v}_i = \widetilde{v}_i$ and $F\bar{v}_\alpha = -\bar{v}_\alpha$. Since $F\xi_1 = \xi_1$ and $F\xi_2 = -\xi_2$, we know that there exist an orthonormal basis $\{\widetilde{w}_1, \dots, \widetilde{w}_{k+1}, \dots, \bar{w}_1, \dots, \bar{w}_{n-k+2}\}$ of B_p, with $\widetilde{g}(\bar{w}_{n-k+2}, \bar{w}_{n-k+2}) = -1$, such that $F\widetilde{w}_i = \widetilde{w}_i$ and $F\bar{w}_\alpha = -\bar{w}_\alpha$. Since $R^D = 0$ and M^n is simply connected, there exist parallel sections $\widetilde{\eta}_1, \dots, \widetilde{\eta}_{k+1}, \bar{\eta}_1, \dots, \bar{\eta}_{n-k+2}$ on B such that $\widetilde{\eta}_i(p) = \widetilde{w}_i$ and $\bar{\eta}_\alpha(p) = \bar{w}_\alpha$. Moreover since the connection D is compatible with the metric $\widetilde{g}$, we have that $\widetilde{\eta}_1, \dots, \widetilde{\eta}_{k+1}, \bar{\eta}_1, \dots, \bar{\eta}_{n-k+2}$ are parallel orthonormal sections on B. From lemma 5, we obtain that $F\widetilde{\eta}_i = \widetilde{\eta}_i$ and $F\bar{\eta}_\alpha = -\bar{\eta}_\alpha$. This completes the proof. □

We are ready to proof theorem 1.

Proof of theorem 1. In Proposition 1 we showed that there exist parallel orthonormal sections on B $\widetilde{\eta}_1, \dots, \widetilde{\eta}_{k+1}, \bar{\eta}_1, \dots, \bar{\eta}_{n-k+2}$ with $k \in \{1, \dots, n\}$ such that $\widetilde{\eta}_1, \dots, \widetilde{\eta}_{k+1} \in B_1$ and that $\bar{\eta}_1, \dots, \bar{\eta}_{n-k+2} \in B_{-1}$ with $k \in \{1, \dots, n\}$. In Lemma 10 we showed that $F^2 = I$ and that F is symmetric with respect to $\widetilde{g}$. Since $\widetilde{\eta}_1, \dots, \bar{\eta}_{n-k+2}$ are parallel orthonormal sections, we know that $\xi_1 = \sum_{i=1}^{k+1} \widetilde{g}(\xi_1, \widetilde{\eta}_i)\widetilde{\eta}_i + \sum_{\alpha=1}^{n-k+2} \epsilon_\alpha \widetilde{g}(\xi_1, \bar{\eta}_\alpha)\bar{\eta}_\alpha$, where $\epsilon_1 = \cdots = \epsilon_{n-k+1} = 1$ and $\epsilon_{n-k+2} = -1$. We obtain that $\widetilde{g}(\bar{\eta}_\alpha, \xi_1) = 0$ for $\alpha \in \{1, \dots, n-k+2\}$, because F is symmetric with respect to $\widetilde{g}$, $F\xi_1 = \xi_1$ and $F\bar{\eta}_\alpha = -\bar{\eta}_\alpha$ and hence $\xi_1 = \sum_{i=1}^{k+1} \widetilde{g}(\xi_1, \widetilde{\eta}_i)\widetilde{\eta}_i$. Analogously, we obtain that $\xi_2 = \sum_{\alpha=1}^{n-k+2} \epsilon_\alpha \widetilde{g}(\xi_2, \bar{\eta}_\alpha)\bar{\eta}_\alpha$. Using the components of ξ_1 and ξ_2 with respect to $\widetilde{\eta}_1, \dots, \widetilde{\eta}_{k+1}, \bar{\eta}_1, \dots, \bar{\eta}_{n-k+2}$ will construct an isometric immersion $\psi : M^n \to \mathbb{S}^k \times \mathbb{H}^{n-k+1}$. Define $\psi : M^n \to \mathbb{L}^{n+3}$ by

$$p \mapsto (\widetilde{g}_p(\widetilde{\eta}_1, \xi_1), \dots, \widetilde{g}_p(\widetilde{\eta}_{k+1}, \xi_1), \widetilde{g}_p(\bar{\eta}_1, \xi_2), \dots, \widetilde{g}_p(\bar{\eta}_{n-k+2}, \xi_2)).$$

It is easy to see that $\sum_{i=1}^{k+1} \widetilde{g}(\widetilde{\eta}_i, \xi_1)^2 = 1$ and $\sum_{\alpha=1}^{n-k+2} \epsilon_\alpha \widetilde{g}(\bar{\eta}_\alpha, \xi_2)^2 = -1$ and hence $\psi(M^n) \subset \mathbb{S}^k \times \mathbb{H}^{n-k+1}$. Next we will show that ψ is an immersion. Let p be an arbitrary point of M^n and v an arbitrary tangent vector at p such that $\psi_* v = 0$. We obtain that

$$0 = v[\widetilde{g}(\widetilde{\eta}_i, \xi_1)] = \widetilde{g}(\widetilde{\eta}_i, D_v\xi_1) = \widetilde{g}(\widetilde{\eta}_i, \frac{1}{2}(v + fv + g(U, v)\xi))$$

and

$$0 = \widetilde{g}(\bar{\eta}_\alpha, \frac{1}{2}(v - fv - g(U, v)\xi)).$$

We also have that

$$\begin{aligned} F(\frac{1}{2}(v + fv + u(v)\xi) &= \frac{1}{2}(fv + f^2 + u(v)U + (g(U, v + fv + \lambda v))\xi) \\ &= \frac{1}{2}(v + fv + u(v)\xi) \end{aligned}$$

and hence

$$\frac{1}{2}(v + fv + u(v)\xi) = 0. \tag{26}$$

Analogously, we obtain that $F(\frac{1}{2}(v - fv - u(v))\xi) = -\frac{1}{2}(v - fv - u(v)\xi))$ and hence

$$\frac{1}{2}(v - fv - u(v)\xi) = 0. \tag{27}$$

Summing up the equations (26) and (27), we obtain that $v = 0$ and hence ψ is an immersion, because p and v were arbitrary. Next we will show that ψ is isometric, i.e. $g(v, w) = \langle \psi_* v, \psi_* w\rangle$, where $\langle ., .\rangle$ is the Lorentzian inproduct on $\mathbb{R}^{n+3}$.

$$\begin{aligned}
\langle \psi_* v, \psi_* w\rangle &= \frac{1}{4}(\sum_{i=1}^{k+1} \widetilde{g}(v + fv + u(v)\xi, \widetilde{\eta}_i)\widetilde{g}(w + fw + u(w)\xi, \widetilde{\eta}_i) \\
&\quad + \sum_{\alpha=1}^{n-k+2} \epsilon_\alpha \widetilde{g}(v - fv - u(v)\xi, \bar{\eta}_\alpha)\widetilde{g}(w - fw - u(w)\xi, \bar{\eta}_\alpha)) \\
&= \frac{1}{4}(g(v + fv + u(v)\xi, w + fw + u(w)\xi) + g(v - fv - u(v)\xi, w - fw - u(w)\xi)) \\
&= \frac{1}{4}(2g(v, w) + 2g(f^2 v, w)2 + 2u(v)u(w)) \\
&= \frac{1}{4}(2g(v, w) + 2g(v, w) - 2u(v)u(w) + 2u(v)u(w)) \\
&= g(v, w).
\end{aligned}$$

We can conclude that ψ is an isometric immersion of M^n into $\mathbb{S}^k \times \mathbb{H}^{n-k+1}$. We will show now that $N = (\widetilde{g}(\widetilde{\eta}_1, \xi), \dots, \widetilde{g}(\bar{\eta}_{n-l+2}, \xi))$ is a unit normal of $\psi(M^n)$ in $\mathbb{S}^k \times \mathbb{H}^{n-k+1}$ and moreover the shape operator of M^n associated to the normal N is given by S. Let p be an arbitrary point of M^n and $v \in T_p M^n$:

$$\begin{aligned}
\langle \psi_* v, N\rangle &= \sum_{i=1}^{k+1} \widetilde{g}(\frac{1}{2}(v + fv + u(v)\xi), \widetilde{\eta}_i)\widetilde{g}(\xi, \widetilde{\eta}_i) \\
&\quad + \sum_{\alpha=1}^{n-k+2} \epsilon_\alpha \widetilde{g}(\frac{1}{2}(v - fv - u(v)), \bar{\eta}_\alpha)\widetilde{g}(\xi, \bar{\eta}_\alpha) \\
&= \widetilde{g}(\frac{1}{2}(v + fv + u(v)\xi), \xi) + \widetilde{g}(\frac{1}{2}(v - fv - u(v)), \xi) \\
&= \frac{1}{2}(u(v) - u(v)) \\
&= 0
\end{aligned}$$

and hence N is a unit normal of $\psi(M^n)$. But we also have that $\langle N, N_1\rangle = \langle N, N_2\rangle = 0$, where $N_1 = (\widetilde{g}(\widetilde{\eta}_1, \xi_1), \dots, \widetilde{g}(\widetilde{\eta}_{k+1}, \xi_1), 0, \dots, 0)$ and $N_2 = (0, \dots, 0, \widetilde{g}(\bar{\eta}_1, \xi_2), \dots, \widetilde{g}(\bar{\eta}_{n-k+2}, \xi_2))$. Therefore N is a unit normal of $\psi(M^n)$ in $\mathbb{S}^k \times \mathbb{H}^{n-k+1}$. Next we show that S is indeed the shape operator of $\psi(M^n)$ in $\mathbb{S}^k \times \mathbb{H}^{n-k+1}$ with respect to N:

$$\begin{aligned}
\langle \psi_* v, \widetilde{\nabla}_{\psi_* v} N\rangle &= \langle \psi_* v, D_{\psi_* v} N\rangle \\
&= \widetilde{g}(\frac{1}{2}(v + fv + u(v)\xi), -Sw) \\
&\quad + \widetilde{g}(\frac{1}{2}(v - fv - u(v)\xi), -Sw) \\
&= g(v, -Sw).
\end{aligned}$$

Suppose now that F is the product structure of $\mathbb{S}^k \times \mathbb{H}^{n-k+1}$. It is easy to deduce by direct calculations that $F(N) = \psi_*(U) + \lambda N$ and that $F(\psi_*(X)) = \psi_*(fX) + u(X)N$. Finally we will prove that the isometric immersion is unique up to a isometry of $\mathbb{S}^k \times \mathbb{H}^{n-k+1}$. Let $\psi_1, \psi_2 : M^n \to \mathbb{S}^k \times \mathbb{H}^{n-k+1}$ be two isometric immersions of M^n in $\mathbb{S}^k \times \mathbb{H}^{n-k+1}$ with unit normals N_1 and N_2, respectively, such that

$$\widetilde{\nabla}_{\psi_{*j}(X)} N_j = -\psi_{*j}(SX) \tag{28}$$

$$F(\psi_{*j}(X)) = \psi_{*j}(fX) + u(X)N_j \tag{29}$$

$$F(N_j) = \psi_{*j}(U) + \lambda N_j, \tag{30}$$

with $j = 1, 2$. We will search now for an isometry ϕ of $\mathbb{S}^k \times \mathbb{H}^{n-k+1}$ such that $\phi \circ \psi_1 = \psi_2$. Define a map $C : M^n \to O^1(n+3)$ by

$$\begin{aligned} C_p(\psi_{*1}(X)) &= \psi_{*2}(X), \\ C_p(N_1) &= N_2, \\ C_p(\widetilde{N}_1) &= \widetilde{N}_2, \\ C_p(\bar{N}_1) &= \bar{N}_2, \end{aligned}$$

for all $X \in T_pM^n$ and whereby $\widetilde{N}_j = (\psi_j^1, \ldots, \psi_j^{k+1}, 0, \ldots, 0)$ and $\bar{N}_j = (0, \ldots, 0, \psi_j^{k+2}, \ldots, \psi_j^{n+3})$, where ψ_j^i are the components of ψ_j with respect to the standard basis of $\mathbb{R}^{n+3}$. We have identified here $T_{\psi_1(p)}\mathbb{R}^{n+3}$ and $T_{\psi_2(p)}\mathbb{R}^{n+3}$ with $\mathbb{R}^{n+3}$. We will show that C is an constant map, by showing that $(D_XC)(V) = D_XC(V) - C(D_XV) = 0$ for every $X \in T_pM^n$ and every vector field V along ψ_1. To prove $(D_XC)(V) = 0$, it is sufficient to consider two cases: V is a tangent vector field along ψ or V is a normal vector field of M^n in $\mathbb{L}^{n+3}$. Assume first that $V = \psi_{*1}(Y)$, where Y is a vector field of M^n. Then

$$\begin{aligned} (D_XC)(\psi_{*1}(Y)) &= D_XC(\psi_{*1}(Y)) - C(D_X\psi_{*1}(Y)) \\ &= D_X\psi_{*2}(Y) - C(D_X\psi_{*1}(Y)) \\ &= \psi_{*2}(\nabla_XY) + g(SX, Y)N_2 \\ &\quad - \frac{1}{2}(g(X, Y + fY))\widetilde{N}_2 + \frac{1}{2}(g(X, Y - fY))\bar{N}_2 \\ &\quad - C(\psi_{*1}(\nabla_XY) + g(SX, Y)N_1 \\ &\quad - \frac{1}{2}(g(X, Y + fY))\widetilde{N}_1 + \frac{1}{2}(g(X, Y - fY))\bar{N}_1) \\ &= 0. \end{aligned}$$

Assume now that $V = N_1$, then

$$\begin{aligned} (D_XC)(N_1) &= D_XC(N_1) - C(D_XN_1) \\ &= -\psi_{*2}(SX) - \frac{1}{2}u(X)(\widetilde{N}_2 + \bar{N}_2) + C(\psi_{*1}(SX) + \frac{1}{2}u(X)(\widetilde{N}_1 + \bar{N}_1)) \\ &= 0. \end{aligned}$$

One can also prove that $(D_XC)(\widetilde{N}_1) = (D_XC)(\bar{N}_1) = 0$, by using $D_X\widetilde{N}_j = \frac{1}{2}(\psi_{*j}(X + fX) + u(X)N_j)$ and $D_X\bar{N}_j = \frac{1}{2}(\psi_{*j}(X - fX) - u(X)N_j)$. Hence we obtain that C is a constant map and can be identified with $\phi \in O^1(n+3)$. Since $\phi(\widetilde{N}_1) = \widetilde{N}_2$ and $\phi(\bar{N}_1) = \bar{N}_2$, we obtain that $\phi(\psi_1) = \psi_2$. □

REFERENCES

[1] B.Y. Chen, Geometry of submanifolds, Marcel Dekker, New York, 1974.

[2] B. Daniel, *Isometric immersions into* $\mathbb{S}^n \times \mathbb{R}$ *and* $\mathbb{H}^n \times \mathbb{R}$ *and applications to minimal surfaces*, Trans. Amer. Math. Soc., to appear.

[3] F. Dillen, *Equivalence theorems in affine differential geometry*, Geom. Dedicata **32** (1989), 81-92.

[4] F. Dillen, K. Nomizu and L. Vrancken, *Conjugate connections and Radon's theorem in affine differential geometry* Monatsh. Math. **109** (1990), 221-235.

Katholieke Universiteit Leuven, Departement Wiskunde, Celestijnenlaan 200 B, B-3001 Leuven, Belgium

E-mail address, D. Kowalczyk: `daniel.kowalczyk@wis.kuleuven.be`